

Laboratory News

VOL. 39, NO. 3 - May 16, 2016

NEW TESTS FOR TACROLIMUS AND CYCLOSPORINE BY LIQUID CHROMATOGRAPHY/TANDEM MASS SPECTROMETRY

Joyce Flanagan, PhD, Clinical Chemist; Jennifer Winterhack, Assistant Manager, Toxicology Lab

Effective May 16th, 2016, Marshfield Labs will offer two new tests for the immunosuppressant drugs tacrolimus and cyclosporine. The tests will use liquid chromatography/tandem mass spectrometry methodology. Tacrolimus and cyclosporine are drugs that inhibit the body's immune response and are typically administered to prevent the rejection of transplanted organs.

TEST INFORMATION

Lab Test Code	Description	CPT Code	Therapeutic Range*	Toxic Level
CYCLO	Cyclosporine	80158	100 - 400 ng/mL	> 700 ng/mL
TACRO	Tacrolimus	80197	5 - 15 ng/mL	> 25 ng/mL

^{*}The optimal therapeutic range for a given patient may differ from the general therapeutic range given above based on type of organ transplant, treatment phase, use in combination with other drugs, time of specimen collection relative to prior dose, and the therapeutic approach of the provider.

CONTACTS

- Clinical questions: Joyce Flanagan; Technical questions: Jennifer Winterhack.
- Phone number: 800-222-5835.

SPECIMEN REQUIREMENTS

Specimen Type: Whole blood only (EDTA or K₂EDTA)

Minimum Volume: 1 mL

Storage/Transport: Refrigerated

Stability: Room temperature: 24 hours;

Refrigerated (4°C): 7 days; Frozen: 30 days.

Rejection Criteria: Serum or plasma; Clotted specimens;

Specimens left at room temp >24 hours.

Available: Mon-Fri once daily; Sample must be in lab by

1:00 PM, result reported by 5:00 PM.

Analytical Time: 1 day 🧌

